

Best Available Copy

GENES

Benjamin Lewin

Oxford New York Tokyo
Oxford University Press
1997

Best Available Copy

Oxford University Press, Great Clarendon Street, Oxford OX2 6DP

Oxford New York

Athens Auckland Bangkok Bogota Bombay Buenos Aires
Calcutta Cape Town Dar es Salaam Delhi Florence Hong Kong
Istanbul Karachi Kuala Lumpur Madras Madrid Melbourne
Mexico City Nairobi Paris Singapore Taipei Tokyo Toronto

and associated companies in
Berlin Ibadan

Oxford is a trade mark of Oxford University Press

Published in the United States
by Oxford University Press, Inc., New York

© Oxford University Press and Cell Press, 1997

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, without the prior permission in writing of Oxford University Press. Within the UK, exceptions are allowed in respect of any fair dealing for the purpose of research or private study, or criticism or review, as permitted under the Copyright, Designs and Patents Act, 1988, or in the case of reprographic reproduction in accordance with the terms of licences issued by the Copyright Licensing Agency. Enquiries concerning reproduction outside those terms and in other countries should be sent to the Rights Department, Oxford University Press, at the address above.

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form of binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

A catalogue record for this book is available from the British Library

Library of Congress Cataloging in Publication Data
(Data available)

Typeset by Wyvern Typesetting Ltd, Bristol

Printed in The United States of America

CHAPTER 27

Nucleosomes

Best Available Copy

Chromatin has a compact organization in which most DNA sequences are structurally inaccessible and functionally inactive. Within this mass are the minority of active sequences. What is the general structure of chromatin, and what is the difference between active and inactive sequences? The high overall packing ratio of the genetic material immediately suggests that DNA cannot be directly packaged into the final structure of chromatin. There must be *hierarchies* of organization.

The fundamental subunit of chromatin has the same type of design in all eukaryotes. The nucleosome contains ~200 bp of DNA, organized by an octamer of small, basic proteins in a bead-like structure. The protein components are histones. They form an interior core; DNA lies on the surface of the particle. Nucleosomes are an invariant component of euchromatin and heterochromatin in the inter-nucleus, and of mitotic chromosomes. Formation of the nucleosome forms the first level of organization, giving a packing ratio of ~6. The components and structure of nucleosome are well characterized.

The second level of organization is the coil-like series of nucleosomes into a helical fiber that constitutes the ~30 nm fiber that is found in both interphase chromatin and mitotic chromosomes (see Figure 26.8). In chromatin, the packing ratio of DNA to ~40. The structure of this fiber requires additional components but is not well defined.

The overall packing ratio is determined by the highest level of organization, the packaging of chromatin itself. This gives an overall packing

ratio of ~1000 in euchromatin, cyclically interchangeable with packing into mitotic chromosomes to achieve an overall ratio of ~10,000. A similar increase in condensation is seen in heterochromatin.

We need to work through these levels of organization to characterize the events involved in cyclical packaging, replication, and transcription. We assume that association with additional proteins, or modifications of existing chromosomal proteins, are involved in changing the structure of chromatin. We do not know the individual targets for controlling cyclical packaging. Both replication and transcription require unwinding of DNA, and thus must involve an unfolding of the structure that allows the relevant enzymes to manipulate the DNA. This is likely to involve changes in all levels of organization, but at present these processes can be characterized only in terms of changes at the level of the nucleosomes.

When chromatin is replicated, the nucleosomes must be reproduced on both daughter duplex molecules. As well as asking how the nucleosome itself is assembled, we must inquire what happens to other proteins present in chromatin. Since replication disrupts the structure of chromatin, it both poses a problem for maintaining regions with specific structure and offers an opportunity to change the structure.

The mass of chromatin contains up to twice as much protein as DNA. Approximately half of the protein mass is accounted for by the nucleosomes. The mass of RNA is less than 10% of the mass of DNA. Much of the RNA consists